




Beyond Fermat's Last Theorem

David Zureick-Brown
Amherst College

Slides available at <http://dmzb.github.io/>

Colby College Colloquium
April 15, 2024

$$a^2 + b^2 = c^2$$



Basic Problem (Solving Diophantine Equations)

Let f_1, \dots, f_m be polynomials with integer coefficients, e.g.,

$$x^2 + y^2 + 1$$

$$x^3 - y^2 - 2$$

$$2y^2 + 17x^4 - 1$$

Basic problem: solve polynomial equations

Describe the set

$$V(f_1, \dots, f_m) = \{(a_1, \dots, a_n) \in \mathbb{Z}^n : \forall i, f_i(a_1, \dots, a_n) = 0\},$$

i.e., the set of integer solutions to those polynomials

Fact

Solving Diophantine equations is difficult.

Basic Problem (Solving Diophantine Equations)

Let f_1, \dots, f_m be polynomials with integer coefficients, e.g.,

$$x^2 + y^2 + 1 = 0$$

$$x^3 - y^2 - 2 = 0$$

$$2y^2 + 17x^4 - 1 = 0$$

Basic problem: solve polynomial equations

Describe the set

$$V(f_1, \dots, f_m) = \{(a_1, \dots, a_n) \in \mathbb{Z}^n : \forall i, f_i(a_1, \dots, a_n) = 0\},$$

i.e., the set of integer solutions to those polynomials

Fact

Solving Diophantine equations is difficult.

Hilbert's Tenth Problem

Theorem (Davis–Putnam–Robinson 1961, Matijasevič 1970)

There does not exist an algorithm solving the following problem:

input: integer polynomials f_1, \dots, f_m in variables x_1, \dots, x_n ;

output: YES / NO according to whether the set of solutions

$$\{(a_1, \dots, a_n) \in \mathbb{Z}^n : \forall i, f_i(a_1, \dots, a_n) = 0\}$$

is non-empty.

This is *known* to be true for many other cases (e.g., $\mathbb{C}, \mathbb{R}, \mathbb{F}_q, \mathbb{Q}_p, \mathbb{C}(t)$).

This is *still unknown* in many other cases (e.g., \mathbb{Q}).

Fermat's Last Theorem - A Marvelous Proof

Theorem (Wiles; Taylor)

For primes $p \geq 3$ the only integer solutions to the equation

$$x^p + y^p = z^p$$

are integer multiples of the triples

$$(0, 0, 0), \quad (\pm 1, \mp 1, 0), \quad \pm(1, 0, 1), \quad \pm(0, 1, 1).$$

Fermat's Last Theorem - A Marvelous Proof

Theorem (Wiles; Taylor)

For primes $p \geq 3$ the only integer solutions to the equation

$$x^p + y^p = z^p$$

are integer multiples of the triples

$$(0, 0, 0), \quad (\pm 1, \mp 1, 0), \quad \pm(1, 0, 1), \quad \pm(0, 1, 1).$$

This took 300 years to prove!

Fermat's Last Theorem - A Marvelous Proof

Theorem (Wiles; Taylor)

For primes $p \geq 3$ the only integer solutions to the equation

$$x^p + y^p = z^p$$

are integer multiples of the triples

$$(0, 0, 0), \quad (\pm 1, \mp 1, 0), \quad \pm(1, 0, 1), \quad \pm(0, 1, 1).$$

This took 300 years to prove!



Fermat's Last Theorem - A Marvelous Proof

Theorem (Wiles; Taylor)



For primes $p \geq 3$ the only integer solutions to the equation

$$x^p + y^p = z^p$$

are integer multiples of the triples

$$(0, 0, 0), \quad (\pm 1, \mp 1, 0), \quad \pm(1, 0, 1), \quad \pm(0, 1, 1).$$

This took 300 years to prove!



Fermat's Last Theorem - A Marvelous Proof

Theorem (Wiles; Taylor)



For primes $p \geq 3$ the only integer solutions to the equation

$$x^p + y^p = z^p$$


are integer multiples of the triples

$$(0, 0, 0), \quad (\pm 1, \mp 1, 0), \quad \pm(1, 0, 1), \quad \pm(0, 1, 1).$$


This took 300 years to prove!




Fermat's Last Theorem - aftermath






Fermat's Last Theorem - aftermath



Fermat's Last Theorem - aftermath





Books



Fermat trolling



Fermat trolling



See <https://youtu.be/ReOQ300AcSU?si=--fAdsPt4HR3N>

Basic Problem: $f_1, \dots, f_m \in \mathbb{Z}[x_1, \dots, x_n]$

Qualitative:

- ▶ Does there **exist** a solution?
- ▶ Do there exist **infinitely many** solutions?
- ▶ Does the set of solutions have some **extra structure** (e.g., geometric structure, group structure).

Basic Problem: $f_1, \dots, f_m \in \mathbb{Z}[x_1, \dots, x_n]$

Qualitative:

- ▶ Does there **exist** a solution?
- ▶ Do there exist **infinitely many** solutions?
- ▶ Does the set of solutions have some **extra structure** (e.g., geometric structure, group structure).

Quantitative

- ▶ How **many** solutions are there?
- ▶ How **large** is the **smallest** solution?
- ▶ How can we explicitly **find** all solutions? (With proof?)

Basic Problem: $f_1, \dots, f_m \in \mathbb{Z}[x_1, \dots, x_n]$

Qualitative:

- ▶ Does there **exist** a solution?
- ▶ Do there exist **infinitely many** solutions?
- ▶ Does the set of solutions have some **extra structure** (e.g., geometric structure, group structure).

Quantitative

- ▶ How **many** solutions are there?
- ▶ How **large** is the **smallest** solution?
- ▶ How can we explicitly **find** all solutions? (With proof?)

Implicit question

- ▶ Why do equations **have** (or fail to have) solutions?
- ▶ Why do some have **many** and some have **none**?
- ▶ What **underlying mathematical structures** control this?

Example: Pythagorean triples

$$3^2 + 4^2 = 5^2$$

$$5^2 + 12^2 = 13^2$$


$$7^2 + 24^2 = 25^2$$

Lemma


The equation

$$x^2 + y^2 = z^2$$

has infinitely many non-zero coprime solutions.



Pythagorean triples



$$\text{Slope} = t = \frac{y}{x+1}$$

$$x = \frac{1-t^2}{1+t^2}$$

$$y = \frac{2t}{1+t^2}$$

Pythagorean triples

Lemma

The solutions to

$$a^2 + b^2 = c^2$$

(with $c \neq 0$) are all multiples of the triples

$$a = 1 - t^2$$

$$b = 2t$$

$$c = 1 + t^2$$

The Mordell Conjecture

Example

The equation $y^2 + x^2 = 1$ has infinitely many solutions.

The Mordell Conjecture

Example

The equation $y^2 + x^2 = 1$ has infinitely many solutions.

Theorem (Faltings)

For $n \geq 5$, the equation

$$y^2 + x^n = 1$$

has only finitely many solutions.

The Mordell Conjecture

Example

The equation $y^2 + x^2 = 1$ has infinitely many solutions.

Theorem (Faltings)

For $n \geq 5$, the equation

$$y^2 + x^n = 1$$

has only finitely many solutions.

Theorem (Faltings)

For $n \geq 5$, the equation

$$y^2 = f(x)$$

has only finitely many solutions if $f(x)$ is **squarefree**, with **degree** > 4 .

Fermat Curves

Question

Why is Fermat's last theorem believable?


- 1 $x^n + y^n - z^n = 0$ looks like a surface (3 variables)
- 2 $x^n + y^n - 1 = 0$ looks like a curve (2 variables)

Mordell Conjecture

Example

$$y^2 = -(x^2 - 1)(x^2 - 2)(x^2 - 3)$$

This is a cross section of a two holed torus.



The **genus** is the number of holes.

Conjecture (Mordell, 1922)

A curve of genus $g \geq 2$ has only finitely many rational solutions.

Fermat Curves

Question

Why is Fermat's last theorem believable?

- 1 $x^n + y^n - z^n = 0$ looks like a surface (3 variables)
- 2 $x^n + y^n - 1 = 0$ looks like a curve (2 variables)
- 3 and has genus

$$(n-1)(n-2)/2$$

which is ≥ 2 iff $n \geq 4$.

Fermat Curves


Question

What if $n = 3$?

- 1 $x^3 + y^3 - 1 = 0$ is a curve of genus $(3 - 1)(3 - 2)/2 = 1$.
- 2 We were lucky; $Ax^3 + By^3 = Cz^3$ can have infinitely many solutions.

Congruent number problem


$$x^2 + y^2 = z^2, xy = 2 \cdot 6$$



$$3^2 + 4^2 = 5^2, 3 \cdot 4 = 2 \cdot 6$$

Congruent number problem

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$



Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?
How many **digits** does the smallest solution have?

Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?
How many **digits** does the smallest solution have?

$$x = \frac{157841 \cdot 4947203 \cdot 52677109576}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441}$$

$$y = \frac{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 157 \cdot 17401 \cdot 46997 \cdot 356441}{157841 \cdot 4947203 \cdot 52677109576}$$

$$z = \frac{20085078913 \cdot 1185369214457 \cdot 942545825502442041907480}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441 \cdot 157841 \cdot 4947203 \cdot 52677109576}$$

Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?

How many **digits** does the smallest solution have?

$$x = \frac{157841 \cdot 4947203 \cdot 52677109576}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441}$$

$$y = \frac{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 157 \cdot 17401 \cdot 46997 \cdot 356441}{157841 \cdot 4947203 \cdot 52677109576}$$

$$z = \frac{20085078913 \cdot 1185369214457 \cdot 942545825502442041907480}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441 \cdot 157841 \cdot 4947203 \cdot 52677109576}$$

The denominator of z has **44 digits!**

Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?

How many **digits** does the smallest solution have?

$$x = \frac{157841 \cdot 4947203 \cdot 52677109576}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441}$$

$$y = \frac{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 157 \cdot 17401 \cdot 46997 \cdot 356441}{157841 \cdot 4947203 \cdot 52677109576}$$

$$z = \frac{20085078913 \cdot 1185369214457 \cdot 942545825502442041907480}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441 \cdot 157841 \cdot 4947203 \cdot 52677109576}$$

The denominator of z has **44 digits**!
How did anyone ever find this solution?

Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?
How many **digits** does the smallest solution have?

$$x = \frac{157841 \cdot 4947203 \cdot 52677109576}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441}$$

$$y = \frac{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 157 \cdot 17401 \cdot 46997 \cdot 356441}{157841 \cdot 4947203 \cdot 52677109576}$$

$$z = \frac{20085078913 \cdot 1185369214457 \cdot 942545825502442041907480}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441 \cdot 157841 \cdot 4947203 \cdot 52677109576}$$

The denominator of z has **44 digits**!
How did anyone ever find this solution?
(Heegner Points)

Assume the Birch–Swinnerton-Dyer conjectures

If you assume \$1,000,000 worth of conjectures, then the equations

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

have **infinitely many** solutions. **How large** is the smallest solution?
How many **digits** does the smallest solution have?

$$x = \frac{157841 \cdot 4947203 \cdot 52677109576}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441}$$

$$y = \frac{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 157 \cdot 17401 \cdot 46997 \cdot 356441}{157841 \cdot 4947203 \cdot 52677109576}$$

$$z = \frac{20085078913 \cdot 1185369214457 \cdot 942545825502442041907480}{2 \cdot 3^2 \cdot 5 \cdot 13 \cdot 17 \cdot 37 \cdot 101 \cdot 17401 \cdot 46997 \cdot 356441 \cdot 157841 \cdot 4947203 \cdot 52677109576}$$

The denominator of z has **44 digits**!
How did anyone ever find this solution?
(Heegner Points)

“Next” solution has **176 digits**!

Back of the envelope calculation (as of 2011)

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

- Num, den(x, y, z) $\leq 10 \sim 10^6$ many, **1 min** on Emory's computers.

Back of the envelope calculation (as of 2011)

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

- Num, den(x, y, z) $\leq 10 \sim 10^6$ many, **1 min** on Emory's computers.
- Num, den(x, y, z) $\leq 10^{44} \sim 10^{264}$ many, **10^{258} mins = 10^{252} years.**

Back of the envelope calculation (as of 2011)


$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

- Num, den(x, y, z) $\leq 10 \sim 10^6$ many, **1 min** on Emory's computers.
- Num, den(x, y, z) $\leq 10^{44} \sim 10^{264}$ many, **10^{258} mins = 10^{252} years.**
- 10^9 many computers in the world – so **10^{243} years**

Back of the envelope calculation (as of 2011)

$$x^2 + y^2 = z^2, xy = 2 \cdot 157$$

- Num, den(x, y, z) $\leq 10 \sim 10^6$ many, **1 min** on Emory's computers.
- Num, den(x, y, z) $\leq 10^{44} \sim 10^{264}$ many, **10^{258} mins = 10^{252} years.**
- 10^9 many computers in the world – so **10^{243} years**
- Expected time until 'heat death' of universe – **10^{100} years.**



Fermat Surfaces

Conjecture


The only solutions to the equation

$$x^n + y^n = z^n + w^n, n \geq 5$$

satisfy $xyzw = 0$ or lie on the lines 'lines' $x = z, y = w$ (and permutations).


The Swinnerton-Dyer K3 surface

$$x^4 + 2y^4 = 1 + 4z^4$$



The Swinnerton-Dyer K3 surface

$$x^4 + 2y^4 = 1 + 4z^4$$



Two 'obvious' solutions – $(\pm 1 : 0 : 0)$.

The Swinnerton-Dyer K3 surface

$$x^4 + 2y^4 = 1 + 4z^4$$

- Two 'obvious' solutions – $(\pm 1 : 0 : 0)$.
- The next smallest solutions are $(\pm \frac{1484801}{1169407}, \pm \frac{1203120}{1169407}, \pm \frac{1157520}{1169407})$.

Problem

Find another solution. (Probably impossible.)

Back of envelope calculation

- 1 **10^{16} years** to find via brute force.
- 2 Age of the universe – **$13.75 \pm .11$ billion years** (roughly 10^{10}).

Sums of cubes

$$1 = 1^3 + 0^3 + 0^3$$

$$2 = 1^3 + 1^3 + 0^3$$

$$3 = 1^3 + 1^3 + 1^3$$

$$3 = 4^3 + 4^3 + (-5)^3$$

$$4 \neq x^3 + y^3 + z^3$$

$$5 \neq x^3 + y^3 + z^3$$

$$6 = 1^3 + 1^3 + 2^3$$

Conjecture (Heath-Brown)

The equation

$$x^3 + y^3 + z^3 = n$$

has an integer solution if and only if n is not 4 or 5 mod 9.

Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 =$$

Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 = 8866128975287528^3 + (-8778405442862239)^3 + (-2736111468807040)^3$$

Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 = 8866128975287528^3 + (-8778405442862239)^3 + (-2736111468807040)^3$$

$$42 = (-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3$$

Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 = 8866128975287528^3 + (-8778405442862239)^3 + (-2736111468807040)^3$$

$$42 = (-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3$$



Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 = 8866128975287528^3 + (-8778405442862239)^3 + (-2736111468807040)^3$$

$$42 = (-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3$$

$$3 = 569936821221962380720^3 + (-569936821113563493509)^3 + (-472715493453327032)^3$$



Solved by Booker–Sutherland

$$32 \neq x^3 + y^3 + z^3$$

$$33 = 8866128975287528^3 + (-8778405442862239)^3 + (-2736111468807040)^3$$

$$42 = (-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3$$

$$3 = 569936821221962380720^3 + (-569936821113563493509)^3 + (-472715493453327032)^3$$

$$114 = x^3 + y^3 + z^3?$$



“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1),$$

“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1),$$

“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ (\pm 71, -17, 2),$$

“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples


$$\begin{aligned} &(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ &(\pm 71, -17, 2), (\pm 2213459, 1414, 65), \quad (\pm 15312283, 9262, 113), \\ &(\pm 21063928, -76271, 17). \end{aligned}$$

“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$\begin{aligned} &(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ &(\pm 71, -17, 2), (\pm 2213459, 1414, 65), \quad (\pm 15312283, 9262, 113), \\ &(\pm 21063928, -76271, 17). \end{aligned}$$




“Generalized” Fermat equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$\begin{aligned} &(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ &(\pm 71, -17, 2), (\pm 2213459, 1414, 65), \quad (\pm 15312283, 9262, 113), \\ &(\pm 21063928, -76271, 17). \end{aligned}$$



Generalized Fermat Equations

Problem

What are the solutions to the equation $x^a + y^b = z^c$?


Generalized Fermat Equations

Problem

What are the solutions to the equation $x^a + y^b = z^c$?

Theorem (Darmon and Granville)

Fix $a, b, c \geq 2$. Then the equation $x^a + y^b = z^c$ has only finitely many coprime integer solutions iff $\chi = \frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 \leq 0$.



Known Solutions to $x^a + y^b = z^c$ with $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} < 1$

$$1^p + 2^3 = 3^2, \quad 2^5 + 7^2 = 3^4$$

$$7^3 + 13^2 = 2^9, \quad 2^7 + 17^3 = 71^2$$

$$3^5 + 11^4 = 122^2$$

$$17^7 + 76271^3 = 21063928^2$$

$$1414^3 + 2213459^2 = 65^7$$

$$9262^3 + 153122832^2 = 113^7$$

$$43^8 + 96222^3 = 30042907^2$$

$$33^8 + 1549034^2 = 15613^3$$

Known Solutions to $x^a + y^b = z^c$ with $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} < 1$

$$\begin{aligned}1^p + 2^3 &= 3^2, & 2^5 + 7^2 &= 3^4 \\7^3 + 13^2 &= 2^9, & 2^7 + 17^3 &= 71^2 \\3^5 + 11^4 &= 122^2 \\17^7 + 76271^3 &= 21063928^2 \\1414^3 + 2213459^2 &= 65^7 \\9262^3 + 153122832^2 &= 113^7 \\43^8 + 96222^3 &= 30042907^2 \\33^8 + 1549034^2 &= 15613^3\end{aligned}$$

Problem (Beal's conjecture)

These are all solutions with $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 < 0$.

Generalized Fermat Equations – Known Solutions

Conjecture (Beal, Granville, Tijdeman–Zagier)

This is a complete list of coprime non-zero solutions such that

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 < 0.$$

Generalized Fermat Equations – Known Solutions

Conjecture (Beal, Granville, Tijdeman–Zagier)

This is a complete list of coprime non-zero solutions such that

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 < 0.$$

\$1,000,000 prize for proof of conjecture...

Generalized Fermat Equations – Known Solutions

Conjecture (Beal, Granville, Tijdeman–Zagier)

This is a complete list of coprime non-zero solutions such that

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 < 0.$$

\$1,000,000 prize for proof of conjecture...

...or even for a counterexample.

Generalized Fermat Equations – Known Solutions

Conjecture (Beal, Granville, Tijdeman–Zagier)

This is a complete list of coprime non-zero solutions such that

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 < 0.$$

\$1,000,000 prize for proof of conjecture...
...or even for a counterexample.

The logo for SETI@home, featuring the text "SETI@home" in white against a dark, starry space background.The logo for ABC@home, featuring the letters "A", "B", and "C" in white on a yellow background with a grid pattern, followed by "@home" in black. Above the "@home" text is the mathematical expression $c < K(\epsilon) \prod_{p|c} p^{1+\epsilon}$.

Examples of Generalized Fermat Equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$\begin{aligned} &(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ &(\pm 71, -17, 2), (\pm 2213459, 1414, 65), \quad (\pm 15312283, 9262, 113), \\ &(\pm 21063928, -76271, 17). \end{aligned}$$

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{7} - 1 = -\frac{1}{42} < 0$$

Examples of Generalized Fermat Equations

Theorem (Poonen, Schaefer, Stoll)

The coprime integer solutions to $x^2 + y^3 = z^7$ are the 16 triples

$$\begin{aligned} &(\pm 1, -1, 0), \quad (\pm 1, 0, 1), \quad \pm(0, 1, 1), \quad (\pm 3, -2, 1), \\ &(\pm 71, -17, 2), (\pm 2213459, 1414, 65), \quad (\pm 15312283, 9262, 113), \\ &(\pm 21063928, -76271, 17). \end{aligned}$$

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{7} - 1 = -\frac{1}{42} < 0$$

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{6} - 1 = 0$$

Examples of Generalized Fermat Equations

Theorem (Darmon, Merel)

Any pairwise coprime solution to the equation

$$x^n + y^n = z^2, n > 4$$

satisfies $xyz = 0$.

$$\frac{1}{n} + \frac{1}{n} + \frac{1}{2} - 1 = \frac{2}{n} - \frac{1}{2} < \frac{2}{4} - \frac{1}{2} = 0$$

Other applications of the modular method

Ideas behind the proof of FLT permeate the study of diophantine problems.

Other applications of the modular method

Ideas behind the proof of FLT permeate the study of diophantine problems.

Theorem (Bugeaud, Mignotte, Siksek; 2006)

The only Fibonacci numbers that are perfect powers are

$$F_1 = F_2 = 1, F_6 = 8, F_{12} = 144.$$

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

Other applications of the modular method

Ideas behind the proof of FLT permeate the study of diophantine problems.

Theorem (Bugeaud, Mignotte, Siksek; 2006)

The only Fibonacci numbers that are perfect powers are

$$F_1 = F_2 = 1, F_6 = 8, F_{12} = 144.$$

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

Theorem (Silliman–Vogt; 2013 REU)

0 and 1 are the only perfect powers in the Lucas sequence

$$L_1 = 0, L_2 = 1, \quad L_n = 3L_{n-1} - 2L_{n-2}.$$

0, 1, 3, 7, 15, 31, 63, 127, 255, 511, 1023, 2047, 4095, 8191, ..., $2^n - 1$, ...

Examples of Generalized Fermat Equations

Theorem (Klein, Zagier, Beukers, Edwards, others)

The equation

$$x^2 + y^3 = z^5$$

Examples of Generalized Fermat Equations

Theorem (Klein, Zagier, Beukers, Edwards, others)

The equation

$$x^2 + y^3 = z^5$$

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{5} - 1 = \frac{1}{30} > 0$$

Examples of Generalized Fermat Equations

Theorem (Klein, Zagier, Beukers, Edwards, others)

The equation

$$x^2 + y^3 = z^5$$

has infinitely many coprime solutions

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{5} - 1 = \frac{1}{30} > 0$$

Examples of Generalized Fermat Equations

Theorem (Klein, Zagier, Beukers, Edwards, others)

The equation

$$x^2 + y^3 = z^5$$

has infinitely many coprime solutions

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{5} - 1 = \frac{1}{30} > 0$$

$$(T/2)^2 + H^3 + (f/12^3)^5$$

- 1 $f = st(t^{10} - 11t^5s^5 - s^{10})$,
- 2 $H = \text{Hessian of } f$,
- 3 $T = \text{a degree 3 covariant of the dodecahedron.}$

(a, b, c) such that $\chi < 0$ and the solutions to $x^a + y^b = z^c$ have been determined.

$\{n, n, n\}$	Wiles, Taylor–Wiles, building on work of many others
$\{2, n, n\}$	Darmon–Merel, others for small n
$\{3, n, n\}$	Darmon–Merel, others for small n
$\{5, 2n, 2n\}$	Bennett
$(2, 4, n)$	Ellenberg, Bruin, Ghioca $n \geq 4$
$(2, n, 4)$	Bennett–Skinner; $n \geq 4$
$\{2, 3, n\}$	Poonen–Shaefer–Stoll, Bruin. $6 \leq n \leq 9$
$\{2, 2\ell, 3\}$	Chen, Dahmen, Siksek; primes $7 < \ell < 1000$ with $\ell \neq 31$
$\{3, 3, n\}$	Bruin; $n = 4, 5$
$\{3, 3, \ell\}$	Kraus; primes $17 \leq \ell \leq 10000$
$(2, 2n, 5)$	Chen $n \geq 3^*$
$(4, 2n, 3)$	Bennett–Chen $n \geq 3$
$(6, 2n, 2)$	Bennett–Chen $n \geq 3$
$(2, 6, n)$	Bennett–Chen $n \geq 3$

(a, b, c) such that $\chi < 0$ and the solutions to $x^a + y^b = z^c$ have been determined.

$\{n, n, n\}$	Wiles, Taylor–Wiles, building on work of many others
$\{2, n, n\}$	Darmon–Merel, others for small n
$\{3, n, n\}$	Darmon–Merel, others for small n
$\{5, 2n, 2n\}$	Bennett
$(2, 4, n)$	Ellenberg, Bruin, Ghioca $n \geq 4$
$(2, n, 4)$	Bennett–Skinner; $n \geq 4$
$\{2, 3, n\}$	Poonen–Shaefer–Stoll, Bruin. $6 \leq n \leq 9$
$\{2, 2\ell, 3\}$	Chen, Dahmen, Siksek; primes $7 < \ell < 1000$ with $\ell \neq 31$
$\{3, 3, n\}$	Bruin; $n = 4, 5$
$\{3, 3, \ell\}$	Kraus; primes $17 \leq \ell \leq 10000$
$(2, 2n, 5)$	Chen $n \geq 3^*$
$(4, 2n, 3)$	Bennett–Chen $n \geq 3$
$(6, 2n, 2)$	Bennett–Chen $n \geq 3$
$(2, 6, n)$	Bennett–Chen $n \geq 3$
$(2, 3, 10)$	ZB